

COMPARATIVE EVALUATION OF THE COMPARABLE SALES METHOD WITH GEOSTATISTICAL VALUATION MODELS

RICHARD BORST
Cole Layer Trumble, US

and

WILLIAM McCLUSKEY
University of Ulster

ABSTRACT

There is an identifiable theoretical relationship between the comparable sales method (CSM) of valuation as practiced by mass appraisers and the recent developments in geostatistical valuation models. This paper provides an evaluation of techniques in respect of their treatment of location and their predictive capability. The CSM is shown to be a special case of a spatially lagged weight matrix model. There is a less formal but clear relationship with Geographically Weighted Regression as well. The predictive accuracy of CSM is compared to several Ordinary Least Squares Model configurations, and results obtained from Geographically Weighted Regression via empirical studies on diverse datasets. An example of a comparable sales weighting scheme as practiced by mass appraisers is provided. In addition, particular interest is focused on how well each method is able to model the spatial variations in property values.

Keywords: Comparable sales, location, geostatistical, spatial autocorrelation, weight matrix, market basket value.

INTRODUCTION

This paper sets out to undertake an empirical comparative analysis of several modeling techniques that specifically address locational characteristics of residential property and how they perform in relation to comparable sales method (CSM). The comparable sales method of valuation as implemented in a mass appraisal setting has gained widespread use in North America, the UK and to a lesser extent Australasia. Given the range of appraisal techniques that can be applied, this research provides for a comparative evaluation to measure whether there is an optimum or best technique that can be employed. However, as to whether it is a “best practice” is a matter for discussion, and even debate. This paper

takes the position that CSM is certainly among the best, if not the best method for the mass appraisal of residential properties. Other candidates for a “best practice” could include the use of a well structured linear or nonlinear model calibrated for an entire jurisdiction, market segmentation models, the “response surface” method, and a collection of advanced modelling techniques. The primary objective of this paper is to compare and evaluate a selected subset of these techniques to the CSM. The main elements being considered in this paper include a description of CSM as implemented in a mass appraisal environment; an introduction to weight matrix models; the relationship of a spatial lag weight matrix model to the comparable sales model; Geographically Weighted Regression (GWR) Models; relationship of GWR to CSM; an empirical comparison of several selected methods; and final concluding comments.

LITERATURE REVIEW OF MASS APPRAISAL TECHNIQUES

Comparable Sales Method

Gipe (1975) recounts the early development history of the automation of the process. An important step in automating the process was the identification of a metric for measuring the comparability of one property to another. That metric has the form

$$D_{jk} = \left(\sum_{i=1}^n [DW_i(X_{ij} - S_{ik})]^2 \right)^{1/2} \quad (1)$$

where D_{jk} is the dissimilarity measure, often referred to as “Distance” between subject property j and sale property k , DW_i is a weighting factor assigned to characteristic i , X_{ij} is the value for the subject property k for the i th variable and S_{ik} is the value of the i th variable for the k th sale. A brief narrative statement of the algorithm will help set the stage for the comparison between comparable sales and the other methods described herein. The main processing steps within the method include finding the n most comparable sales properties (comps); computing an adjusted sale price for each comp; weighting these estimates according to their similarity to the subject; and summing the weighted comp estimates to get the final estimate

Consider that finding the most comparable sale is equivalent to finding the least dissimilar. The actual dissimilarity measure can be based on physical separation, and differences in physical characteristics, date of sale, and the neighborhood to which the comp belongs. The weights in equation (1) can be determined in a number of ways. An example is given in the section entitled “CSM”.

Initially consider estimating the value of a subject based on one comparable sale. Cannaday (1989) describes the method of adjustment of a comparable sale by using the difference in the MRA estimates of the subject and the comparable. This adjustment process can be expressed as:

$$\text{Estimate of } S_k \text{ based on } C_j = ASP(C_j) + (ESP(S_k) - ESP(C_j)) \quad (2)$$

where S_k refers to subject k , C_j to comparable sale j , $ASP(C_j)$ is the actual selling price of C_j , and $ESP(S_k)$ refers to the estimated selling price of subject k . Equation (2) can be rewritten as:

$$\text{Estimate of } S_k \text{ based on } C_j = ESP(S_k) + (ASP(C_j) - ESP(C_j)) \quad (3)$$

where the expression $(ASP(C_j) - ESP(C_j))$ is the residual error of the estimate for comparable sale j . Generalizing (3) to the case of valuing a subject by using several comparable sales results in:

$$CSM(S_k) = \sum_{j=1}^n CW_{jk} [ESP(S_k) + (ASP(C_j) - ESP(C_j))] \quad (4)$$

$$\text{with } \sum_{j=1}^n CW_{jk} = 1$$

where $CSM(S_k)$ refers to the comparable sales method, CW_{jk} is a “comparability weight applied to each comparable sale property. Noting that $\sum_{j=1}^n CW_j ESP(S_k) = ESP(S_k)$, (4) can be rewritten as:

$$CSM(S_k) = ESP(S_k) + \sum_{j=1}^n CW_{jk} (ASP(C_j) - ESP(C_j)) \quad (5)$$

In matrix notation, this expression can be rewritten as:

$$CSM(S) = X \hat{\beta} + CW(Y - X \hat{\beta}) \quad (6)$$

where $\hat{\beta}$ are the estimates of β obtained by OLS methods. The formulation of (6) can be rewritten in the compact form:

$$\tilde{Y} = \hat{Y} + CW \hat{\varepsilon}$$

where the substitution of variables is $CSM(S) = \tilde{Y}$, $\hat{Y} = X \hat{\beta}$, and $W \hat{\varepsilon}$ represents the weighted residual errors from OLS. However, it is the formulation of (6) that will be useful in the comparison to the weight matrix methods.

Thompson (2006) describes a weighting mechanism in common use for developing CW_{jk} combining the n comparable sales estimates expressed as

$$CW_{jk} = 1 / [(D_{\max} / 2)^2 + D_{jk}^2 + (2D_{\max} P_{jk})^2]$$

where:

CW_{jk} = comparability weight of j th sale to the k th subject

D_{\max} = maximum acceptable comparability distance

D_{jk} = actual comparability distance between j th sale and k th subject

P_{jk} = fractional adjustment to the j th sale for the k th subject

Table 1 illustrates the method for $n = 5$.

Table 1: Example of comparable weights computation

Sale Number	1	2	3	4	5
Price	45,000	30,000	50,000	25,000	40,000
Adjusted Selling Price	45,000	40,000	50,000	40,000	30,000
Adjustment to Price	0	10,000	0	15,000	-10,000
P_i	0	0.333	0.000	0.600	-0.250
D_i	10	60	70	80	120
$(D_{\max} / 2)^2$ $D_{\max} = 100$	2,500	2,500	2,500	2,500	2,500
D_i^2	100	3,600	4,900	6,400	14,400
$(2D_{\max} P_i)^2$	0	4444	0	14400	2500
Total Denominator	2,600	10,544	7,400	23,300	19,400
$W_i \times 10,000$ $\sum W_i \times 10,000 = 7.091$	3.846	0.948	1.351	0.429	0.515
Normalized W_i	0.542	0.134	0.191	0.061	0.073
Weighted Contribution	24,410	5,350	9,529	2,421	2,181
Weighted Estimate	43,891				

This methodology is in widespread use in North America. Representative references to its use are found in Underwood and Moesch (1982), Thompson and Gordon (1987),

McCluskey and Borst (1998) and Todora and Whiterell (2002). Additional analytical foundations for the method are described in Colwell et al (1983). Vandell (1991) provides the theoretical background to provide decisions about how many comparables to select, what the comparable selection criteria should be, and how proper weights for each adjusted value estimate can be determined such that the final value estimate is both unbiased and of minimum variance. Kang and Reichert (1991) compared MRA to the comparable sales method on data sets with a variety of characteristics. Gau et al (1992) extended Vandell's method. Lai and Wang (1996) compare the accuracy of the minimum variance grid method to MRA value estimates. They establish the theoretical basis upon which it is shown that the grid method should have a lower variance than MRA estimates.

Weight Matrix Model

In this discussion, W is an $n \times n$ matrix, the elements of which represent the “strength” of the connections between each property and all other properties. There are two variants of the weight matrix model. In one, the matrix is used to model a spatial process directly. These models are referred to as spatial lag models. The other uses a weight matrix to model the error term. Only the former is addressed herein because it has a clear relationship to the comparable sales method.

The simplest form of the spatially lagged weight matrix model is also referred to as the autoregressive model, meaning that the dependent variable is related to itself in a specific way. The logic behind the autoregressive model is that there are “spill over” effects in which the sales prices of nearby properties affect the value of a given property more than those that are farther away. Stated another way, values at close-by locations are more correlated than values at locations that are far apart. This autoregressive model has the following formulation:

$$Y = \rho WY + \varepsilon$$

where ρ is the coefficient of autocorrelation variable, W is a matrix of weights with elements W_{ij} that specify the strength of the relationship between properties i and j , and ε is the error term. In other words, the value of a single observation of the dependent variable, Y_i , is a weighted average of its neighbouring properties Y_j (with $i \neq j$). Models of this form are often referred to as lattice models because the weight matrix is developed for a specific set of points which can be from a regular or irregular arrangement of the points on the two-dimensional plane. Specification of spatially autoregressive model in hedonic form was reported by Can (1990, 1992). The model is expressed as:

$$Y = X\beta + \rho WY + \varepsilon$$

It is often referred to as a mixed regressive, autoregressive model because it is a combination of the basic hedonic model and the autoregressive model forms. Pace and Gilley (1997) use a somewhat different model specification of a weight matrix spatially autoregressive model. In the equation:

$$Y = X\beta + \rho W(Y - X\beta) + \varepsilon$$

the term $\rho W(Y - X\beta)$ represents a weighted average of the errors on nearby properties. As before, W is a matrix of weights W_{ij} with the added properties that $W_{ii} = 0$, the rows of W sum to 1, $0 \leq \rho < 1$ and ε is distributed $N(0, \sigma^2 I)$. This specification is referred to as a simultaneous autoregressive specification (SARS) model with log-likelihood function:

$$L(\rho, \beta, \sigma^2) = \frac{1}{2} \ln |B| - \frac{1}{2} \left[n \ln(2\pi\sigma^2) + \sigma^{-2} (Y - X\beta)' B (Y - X\beta) \right] \quad (7)$$

where $B = (I - \rho W)'(I - \rho W)$. The maximum likelihood (ML) estimation method efficiently estimates the model asymptotically provided that the model assumptions are true. The study in the cited reference compares OLS to SARS on an often cited set of data originally presented by Harrison and Rubinfeld (1978). The SARS method has a lower (44% reduction) error and more intuitively satisfying coefficients.

Pace and Gilley (1998) show how a linear combination of an OLS estimate and comparable sales estimate form the basis of a simultaneous autoregressive (SAR) model. The parameters of this model were estimated by two methods, Estimated Generalized Least Squares (EGLS) and Maximum Likelihood (ML) as mentioned above. A comparison of the two estimating methods was performed on the same sample data used by Can (1992). It was found that the EGLS SARS model was superior in performance relative to OLS and the Grid estimator. In the study of 563 properties, the OLS had 27.9% and grid estimator had 11.7% greater median absolute error than the EGLS SARS. They observe that although the EGLS SARS is superior, the grid estimator provided evidence of its utility in the presence of spatial information. It was called a “poor man’s” spatial autoregression.

Finding solutions by ML involves finding the log determinant of a matrix. When the number of observations is large, and the weighting function is such that the weights are non-zero for a relatively small number of observations in the vicinity of a specific observation, the matrices become large ($N \times N$). Direct methods for finding the log determinant become computationally intense. Barry and Pace (1999) provide a Monte

Carlo simulation method that has been shown to be efficient on matrices of size 1,000,000 x 1,000,000. Such techniques are a necessity for ML techniques to be useful for prediction applications such as assessment. Kelejian and Prucha (1998) describe the theoretical background for a generalized spatial two-stage least squares procedure for estimating a model of the type described in this section. They state that it is computationally simple procedure for estimating models with both a spatially lagged independent variables and error terms. It is fair to say that calibration of such models goes well beyond the skills, expertise and software tools needed to calibrate OLS models.

Relationship of Comparable Sales Model to Weight Matrix Model

Based on the presentation in Pace and Gilley (1998), consider the following combination of the comparable sales estimator and the OLS estimator:

$$\tilde{Y} = \alpha X \hat{\beta} + (1 - \alpha)[X \hat{\beta} + W(Y - X \hat{\beta})] \quad (8)$$

this can be rewritten as:

$$\tilde{Y} = X \hat{\beta} + (1 - \alpha)[W(Y - X \hat{\beta})]$$

or

$$\tilde{Y} = X \hat{\beta} + \rho^*[W(Y - X \hat{\beta})]$$

The analogy to equation (4) is evident. It is presented again here for ease of comparison:

$$Y = X \beta + \rho W(Y - X \beta) + \varepsilon \quad (9)$$

The difference being that the parameters of (9) have yet to be estimated and require advanced calibration methods, while those in (8) could have been estimated by OLS with systematic variation in ρ^* providing an estimate of its optimum value.


Geographically Weighted Regression

In a typical application of Multiple Regression Analysis (MRA), one equation is calibrated on a given set of sales, each of which is weighted equally. There are variants of this technique that weight the points in the sample data set differently, such as Robust Regression, which seeks to down-weight or eliminate outliers in the data set. However, there is still only one equation developed for the entire data set, and there is no inherent spatial component to the weighting method. GWR, on the other hand, is a computationally intensive technique that weights each point in the dataset based on its location. The introduction to the concepts involved in GWR often includes description of moving window regression. In this case, the sample points within a fixed distance of a

given point are included in the regression, all with equal weight, and all others are excluded. This was the case with Dubin's procedure in Case et al (2004). She took the 200-300 points nearest the sample point and included them in the regression model. This procedure was repeated for each data point in the sample set. GWR operates in a similar fashion; however the points do not receive equal weight. Instead, the weight is a function of location, and diminishes with the distance from the regression point.

Figure 1 illustrates the concept of a weighting function based on the coordinates of the regression point and the data points near it. The peak of the surface is the regression point; any sample points under the surface would receive the weight based on the height of the surface at that point.

Figure 1: Example of Spatial Weighting Kernel



Two dimensional representations of the kernel density function are used to illustrate the concept of fixed and variable bandwidth weighting. Figure 2 shows the spatial dimension along the X axis, and the weighting function is represented by the vertical height of the curve. The height of the curve at point j, given by w_{ij} , is the weight applied to point j when point i is the regression point and d_{ij} is the distance between the regression point i and data point j. The bandwidth of the spatial kernel, h , is a parameter that affects how the weight is computed as the distance between the regression point and other sample points increases. It can be fixed or variable. This becomes an important consideration when the sample points are not regularly spaced. A fixed bandwidth could result in there being insufficient points considered in the regression if the bandwidth is too small.

Figure 2: Spatial Kernel





Figure 3 shows two kernels. In the left part of the figure, the ten data points are more closely spaced than the ten points on the right. In this hypothetical example, it can be seen that by appropriately adjusting the bandwidth, ten data points can be considered in each case.

Figure 3: Variable Bandwidth Kernels



The adaptive kernel is more appropriate for application with real property transaction data. This can be reasoned by consideration of the variations in the spatial density of property locations. The number of points contained in a fixed bandwidth does vary

considerably in all three of the counties of the study data, thus all GWR estimation reported herein was performed with an adaptive bandwidth kernel.

Relationship of GWR to Comp Sales

The relationship between GWR and CSM lends itself better to a narrative description. In GWR, a localized model is developed by weighting points based on their distance from the regression point. More distant points receive less weight than those that are closely proximal to the regression point. The regression process itself accounts for the differences in the independent variables for each property included in the regression. CSM achieves its “localization” in two ways. First, physical distance can be incorporated into the dissimilarity measure. This places emphasis on finding nearby properties. However, it also allows for weighting on physical characteristics as well. This allows for finding properties more like the subject, for example in story height, living area, and number of bedrooms. The choice is only limited by the characteristics available in the database of property descriptors. Furthermore, the variables used in the selection process need not have been used in the OLS model used to adjust the comparable sales weights.

Response Surface Methods

The term “Location Value Response Surface” (LVRS) emerged from the assessment community starting in the early 1980’s. Most often, it refers to the application of a location based correction factor to a base value determined via multiple regression analysis or similar multivariate technique without explicit incorporation of location in the model structure. In one of the first reports of its use, O’Connor (1982) described a methodology for developing a location based correction that was in turn applied to a multiplicative form of a valuation model. A model using various functions of x, y coordinates including the distance to certain “Value Influence Centres” (VICS) as independent variables, and the relative value of a typical home at 86 points in the study area was developed as the “Response Surface”. A separate model using building and land characteristics as independent variables and selling price as the dependent variable in a multiplicative form was developed to arrive at a to-be-corrected value estimate. A final value estimate is the product of the base value and the response surface correction computed for the exact location of the property being valued. The LVRS technique evolved over time. O’Connor and Eichenbaum (1988) and Eichenbaum (1988) provide additional insight as to how the VICS are incorporated into the model.

By 1999, the terminology for some authors in the assessment community had changed from LVRS to Response Surface Analysis (RSA) and GIS technology had advanced in capability and accessibility. That is, its migration to desktop computers had made it available to a larger number of users. Ward et al (1999) detailed the use of GIS to develop a surface of normalized sale price per square foot of living area. The normalized factor derived from the surface was utilized as an independent variable in the CAMA model. McCluskey et al (2000) investigate alternative approaches which specifically model the spatial distribution of house prices with the objective of developing location adjustment factors. These approaches were based on the development of surface response

techniques such as inverse distance weighting and universal Kriging. The results generated from the surfaces created were then calibrated within a model structure including other descriptive detail about the sale properties. Ward et al (2002) provide an extensive study of the use of Global Response Surface (GRSA) tools to develop location factors for use as an independent variable in a nonlinear model formulation. Instead of using price per square foot (meter) or price as the dependent variable in the response surface, the concept of a “z-score” ($z = (x - \mu) / \sigma$) which allows for combining data having different fundamental means (μ) and standard deviations (σ) distributions.

METHODOLOGY

The primary methodology adopted in this paper involves the application of each of the techniques to three datasets of residential property. The objective is to evaluate the predictive power of each of the techniques and through a comparison of the coefficient of dispersion (COD) draw conclusions as to the most appropriate for the mass appraisal of residential property. The COD is an internationally recognised statistical measure of model predictive accuracy. In addition, for a subset of the models evaluated, we also presents the results in a format suggested by Thibodeau (2003) which is based on a standardised percentage of properties valued within levels of accuracy.

In this paper, we compare the following model/methods on datasets from three U.S. counties. The primary objective is to test the predictive power of the different methods in estimating sale price:

- OLS regression – baseline global model
- OLS regression – segmented models
- OLS regression – segments as binary variables in global model
- A response surface model
- A combined segmented and response surface model
- GWR regression
- CSM

The Study Data

Sales transaction and property descriptive data have been acquired from three rather different counties in the US. They are Catawba County, North Carolina, Sarasota County, Florida, and Fairfax County, Virginia. Table 2 provides a high level summary of the three jurisdictions. Catawba County is a relatively small, low population density jurisdiction. Sarasota County has a higher population density, but of importance is the much larger amount of “water area”. The number of persons per household is much lower than either Catawba or Fairfax indicative of an older population. This is consistent with the theme of Florida as a place to retire. It has a coastline on the Gulf of Mexico, and considerable inland water resources. It has more than triple the number of housing units of Catawba.

Fairfax County is a high population density suburban community bordering on Washington, DC. It has a much higher average income and more than six times as many housing units than does Catawba.

Table 2: Brief Comparison of Subject Counties

County	State	Pop (1990)	Housing Units	Land Area sq KM	Water Area sq KM	Pop Density	Per Hshld
Catawba	North Carolina	118,412	49,192	1036	35.069	114.302	2.4071
Sarasota	Florida	277,776	157,055	1481	397.61	187.58	1.7687
Fairfax	Virginia	818,584	307,966	1025	29.109	798.976	2.658

There are differences in the housing stock as well. The major differences are shown in Table 3. Each data set was pared down from the original files obtained from the jurisdiction. Some reasonability checks on the relation between the selling price and the property characteristics were employed as well restricting the study to exclude condominiums and town homes.

Table 3: Summary Statistics by County

Variable	County	Count	Mean	St. Dev	Std. Error	Minimum	Maximum	Range
Price	Catawba	7107	159,941	107,103	1270	22,000	1,800,000	1,778,000
	Sarasota	25276	295,036	358,283	2254	13,000	11,000,000	10,987,000
	Fairfax	21700	648,346	317,537	2156	175,000	11,700,000	11,525,000
Sale Date	Catawba	7107	Feb-03			Jan-00	Aug-05	71 months
	Sarasota	25276	Nov-03			Jan-02	Oct-05	43 months
	Fairfax	21700	Jan-05			Jan-04	Jan-06	25 months
Living Area	Catawba	7107	1737	764	8.97	384	8024	7640
	Sarasota	25276	1730	737	4.64	260	10867	10607
	Fairfax	21700	2193	1096	7.44	448	10341	9893
Lot Size (Sq. Ft.)	Catawba	7107	33,343	93,357	1096	3485	3,858,109	3,854,624
	Sarasota	25276	16,699	77,578	488	1283	10,890,000	10,888,720
	Fairfax	21700	19,407	36,055	245	871	2,583,979	2,583,108
Age	Catawba	7107	25.8	23.45	0.275	1	187	186
	Sarasota	25276	25.8	16.64	0.105	2	110	108
	Fairfax	21700	30.4	17.57	0.113	1	136	135

There are several observations about the data that highlight the differences in the housing stock of the three counties:

- Sarasota has a mean selling price nearly double that of Catawba, while Fairfax is approximately four times as high.
- The selling prices start at \$175,000 in Fairfax
- The range of prices for both Sarasota and Fairfax is over six times as large as that of Catawba.
- The sale date range is quite different among the counties
- The homes of Fairfax are somewhat older and larger than that of the other two counties.
- The lot sizes in Catawba are considerably larger than those of the other two counties, consistent with the lower population density.

Histograms of “Price” illustrate the price differential among the three counties (see Figures 4, 5 and 6). All three histograms were trimmed at \$1,000,000 to make the comparison more direct.

Figure 4: Catawba County Sales




Figure 5: Sarasota County Sales





Figure 6: Fairfax County Sales



The variables available for use in the model specification and calibration process are somewhat different in each county. The one thing in common is that they are obtained from an assessment database, and are quite rich in information.

OLS Models- Baseline global model

Establishing a baseline global model for each county is a multi-step process. First, the initial data for each county needs to be transformed into a set of variables suitable for calibrating a model. Once the data have been transformed, a suitable model structure needs to be selected. The global model is based on structural and land characteristics and does not contain specific location information. Similarly, the segmented models contain no explicit location information. The segmentation process is the means by which locational effects are addressed. Three model structures were evaluated for each county – linear, semi-log and log-linear. The model structure with the best performance was chosen individually by county.


OLS Models with Binary Market Segments

An alternative use of the segments identified by the GWR process is to extend the baseline model by incorporating them as binary (dummy) variables. There are several possibilities for the outcome of such models. If their performance is equal to or superior to that of the segmented models, it could be concluded that all of the spatial variation can be accounted for in one model, thus obviating the need for segmented models. This would reduce the segmentation process to one of finding useful “super neighbourhoods”. If, on the other hand, the performance of the extended baseline model is inferior to the segmented model, then further support for the fundamental need for market segmentation has been provided

GWR Models

In brief, a GWR model is developed for each of the three counties. That model is used to value a “market basket value” (MBV) home at each of the regression points in the data. This allows for the formation of a surface, the variation of which represents the composite change in the calibrated model across the study area. Figure 7 provides an example surface. Submarkets are said to exist when the equation describing a candidate submarket is different from a reference equation. This principle is used in forming the submarkets.

Figure 7: Fairfax County MBV Surface




Goodness of Variance Fit optimization, Smith (1986), is used to partition the MBV's into submarkets. The optimal number of submarkets is found by a variety of techniques including COD minimization, Akaike Information Criterion minimization and spatial autocorrelation in residual errors minimization. Here we present the results for the global model and for the best segmented market models.

GWR modelling is a computationally intense process. Processing times increase dramatically with the number of points in the dataset and the number of independent variables considered in the regression. To make the best comparison considering this limitation, all data points were used in each calibration run, but the number of independent

variables were chosen based on the principle of parsimony. Multivariate variable selection routines were used to select the number of variables. With reference to Figure 8, it can be seen that there is little change in R-Squared after ten variables enter the model. Thus, in the case of Catawba county, ten variables were used in GWR.

Figure 8: Variable Selection Curve - Catawba County



Response Surface Models

Ward et al (2002) used the z-score surface as an independent variable in an OLS model formulation. Instead of a z-score, we use the MBV surface value normalized to its mean value as an independent variable. The fundamental difference is in the fact that the MBV is based on all the information in the independent variables.

Combined Segmented and Response Surface Models

All segmented models were recalibrated using the MBV factor as an additional independent variable in the model.

CSM

The procedure outlined in the section, Comparable Sales Method, was utilized to compute values on all properties in each of the three counties. The number of comparable properties considered was set to five. The selection of comparable weights is usually done from both statistical and aesthetic viewpoints. An example will illustrate. In this example, eleven variables were considered in the comparability weighting scheme. A

certain penalty weight is first selected for each variable. In general, it is preferable to have the comparables drawn from the same neighborhood. They are more likely to share the same market influences, and are also more likely to be appreciated by a non-appraiser viewing the evidence supporting the value. As a starting point, a weight of 150 points is set as the “penalty” for leaving a neighborhood. The weight for the remaining variables is set within this context. As an example, the question is asked how much smaller or larger a house can be in a given neighborhood before it would be preferable to look outside the neighborhood. A representative set of weights is provided in Table 4. The last variable in the list is Parcel ID. A very large weight is placed on this variable to insure that the subject property (all of which are also sale properties) is not included as its own comparable. This would give an artificially low error in the predicted value.

Table 4: Representative Set of Weights

Variable Name	Type	Weight
NBHD Group	Binary	250
NBHD	Binary	150
Reverse Month of Sale	Continuous	1
Story Height	Continuous	10
Number of Beds	Continuous	15
Age	Continuous	1
Pool Area	Continuous	0.05
Garage Area	Continuous	0.05
Xcoord	Continuous	0.01
Ycoord	Continuous	0.01
SFLA	Continuous	0.1
PARID	Special	1000

Model Performance Results

The results of the studies described above are summarized in the following tables and charts. Note that the global (baseline) models had no location influence factors in the model structure. They are based on physical characteristics and land size. The relatively high CODs are not presented as representative of a good modelling effort, but rather they serve as a baseline of comparison for the methods described herein to incorporate location into the model. They are simply reference points to show the improvement to be gained by the several methods described herein. The following summary of the methods tested are restated for convenience of reference: The numbering of each item is cross referenced in the table headings.


1. Global OLS Model of best form
 - a. Linear – Catawba (see Exhibit 1)
 - b. Log-linear – “best” in Fairfax and Sarasota (see Exhibit 2 and 3)
2. Segmented Model with Best Number of Segments derived via market basket value from GWR regression surface. In all three cases ten segments were used.
3. Best Global Model using Market Segments as Binary Variables
4. Global Model with normalized Market Basket Value as an Independent Variable
5. Best Segmented Model with normalized Market Basket Value as Independent Variable
6. Global GWR Model – same form as OLS models
7. Comp Sales Method using best predictions from segmented model as adjustment factors

Table 5 provides the performance of each model as measured by the Coefficient of Dispersion. Figures 9 and 10 present the same information in graphical form. The figures use a ratio of each model’s performance to the baseline COD.

Table 5: Comparison of COD Among All Tested Methods

	Number of Sales in Model	Best Global Model	Best Segmented Model	Best Binary Model	MBV as Independent Variable in Global Model	MBV and Best Segmented Model	GWR Global	Comps Method
		(1)	(2)	(3)	(4)	(5)	(6)	(7)
Catawba	7,107	11.05%	10.21%	11.00%	11.08%	10.55%	9.97%	9.75%
Fairfax	19,983	8.94%	6.95%	7.22%	7.05%	6.88%	7.30%	6.52%
Sarasota	24,616	18.99%	14.36%	15.68%	16.09%	14.09%	13.15%	12.92%

Figure 9: Relative Performance of the Methods by County by Model Type



The results of the analysis contained in Table 5 illustrate that the Comparable Sales Method produces the lowest COD for each of the datasets. The importance of these results indicate that whilst improvements are being made in refining surface analysis and segmented location techniques, the CSM provides superior results. The CSM is a more intuitive technique and to a large extent requires considerable less time to produce results, for example, in comparison to Geographically Weighted Regression. The second best performing technique was not as clear. In two of the counties, the GWR model was second in terms of prediction accuracy. In one, Fairfax, the segmented model with a location factor included as an independent variable was second best. We note that the GWR method is computationally intense and is two orders of magnitude slower than traditional OLS techniques. This is not viewed as a major disadvantage because it is expected that the performance of GWR will improve over time with advances in software and hardware.

The performance measure suggested by Thibodeau (2003) is based on the percentage of estimated values that fall within 10% of the transaction value. The standard he puts forth is that a commercial automated valuation system should achieve at least 50% of the estimates within 10% of the sale price. We have expanded that analysis to include additional accuracy ranges as shown in Table 6. These results are consistent with the

patterns found when using COD as the performance measure. Namely, segmented models are more accurate than the baseline model, and that the optimal comparable sales method is the most accurate. Catawba and Fairfax achieve compliance with the minimum criteria starting with the baseline model, and improving with segmentation and the use of comparable sales. In Sarasota, the accuracy measure is only achieved with the optimal comparable sales method.

Table 6: Comparison of Accuracy of the Three Models

	Catawba			Fairfax			Sarasota		
Accuracy Range	Baseline	Segmented	Optimal Comps	Baseline	Segmented	Optimal Comps	Baseline	Segmented	Optimal Comps
95-105%	32.01%	35.22%	38.19%	37.40%	47.52%	51.26%	18.38%	26.73%	30.54%
90-110%	58.60%	61.83%	65.19%	65.05%	77.29%	80.87%	35.42%	49.51%	55.39%
85-115%	75.97%	78.47%	79.81%	82.36%	90.44%	92.24%	50.51%	66.25%	71.44%
50-150%	98.68%	99.20%	99.27%	99.75%	99.89%	99.90%	93.13%	96.73%	97.25%

CONCLUSIONS

This paper set out to undertake an empirical comparative analysis of several modeling approaches that specifically address locational characteristics of real property and how they perform in relation to comparable sales method (CSM). The use of comparable sales is an intuitive valuation technique and used in conjunction with multiple regression analysis, it produces high levels of predictive accuracy. The importance of the location factor in deterministic residential appraisal models has resulted in the development of techniques which are spatial and/or geostatistically based. These contemporary techniques would appear to have promise in terms of more accurately predicting residential property values. The common theme within all of the techniques is the identification of ‘neighbourhoods’ or ‘locational segments’.

Historically, the CSM has been considered as being the pre-eminent methodology for the mass appraisal of residential property. Given the development of new innovative approaches, it is important to be able to assess their capability in terms of predictive accuracy. Those working within the mass appraisal environment provide testament to their needs for research and development in terms of delivering workable techniques. In view of this, the research has attempted to comparatively evaluate a number of the more recent modeling alternatives with CSM. The results of this research show that the traditional comparable sales method has been demonstrated to be superior in predictive performance to all other methods presented. The results of CSM are significant when measured against the performance standard as suggested by Thibodeau (2003). In addition, these results are consistent with the patterns found when using COD as the performance measure.

Namely, segmented models are more accurate than the baseline model, and that the optimal comparable sales method is the most accurate. Clearly, this has implications for the mass appraisal community in terms of the development of statistically sound and accurate models.

Whilst these study areas are confined to the United States, they represent a wide variety of market conditions typically to be found in many jurisdictions and not unique to the US. Thus, the methodology evaluated is believed to be applicable to any region of the world having open market transactions of real estate on domestic properties.

A further conclusion of this research relates to the fact that the future development of mass appraisal techniques will lie in the direction of hybrid interactive models. It is essentially the marrying together of traditional approaches such as regression with spatially derived elements resulting in a technique such as geographically weighted regression. Whilst this may be the future direction such developments require industry acceptance and support if they are to flourish as the 'new' techniques of the twenty first century.

REFERENCES

- Barry, R.P. and Pace, R.K. (1999), Monte Carlo Estimates of the Log Determinant of Large Sparse Matrices, *Linear Algebra and its Applications*, 289, 41-54.
- Can, A. (1992), Specification and Estimation of Hedonic Housing Price Models, *Regional Science and Urban Economics*, 22, 453-474.
- Cannaday, R.E., (1989), How Should You Estimate and Provide Market Support for Adjustments in Single Family Appraisals, *Real Estate Appraiser and Analyst*, 55:4, 43-54.
- Case, B., Clapp, J., Dubin, R. and Rodriguez, M. (2004), Modeling Spatial and Temporal House Price Patterns: A Comparison of Four Models, *Journal of Real Estate Finance and Economics*, 29:2, 167-191.
- Colwell, P.F., Cannaday, R.E. and Wu, C. (1983), The Analytical Foundations of Adjustment Grid Methods, *Journal of the American Real Estate and Urban Economics Association*, 11:1, 11-29.
- Eichenbaum, J. (1988), Incorporating Location into Computer Assisted Valuation: Progress in New York City, paper presented at the World Congress III Computer-Assisted Valuation and Land Information Systems Assisted Valuation, Cambridge, MA.
- Gau, G.W., Lai, T-Y and Wang, K. (1992), Optimal Comparable Selection and Weighting in Real Property Valuation: An Extension, *AREUEA Journal*, 20:1, 107-123.
- Gipe, G.W. (1974), Analysis of Residuals to Improve Multiple Regression Equations, *Assessors Journal*, 9:1, 9-16.

- Harrison, D. and Rubinfeld, D.L. (1978), Hedonic Housing Prices and the Demand for Clean Air, *Journal of Environmental Economics and Management*, 5, 81-102.
- Kang, H-B. and Reichert, A.K. (1991), An Empirical Analysis of Hedonic Regression and Grid-Adjustment Techniques in Real Estate Appraisal, *AREUEA Journal*, 19:1 70-91.
- Kelejian, H.H. and Prucha, I.R., (1998), A Generalized Spatial Two-Stage Least Squares Procedure for Estimating a Spatial Autoregressive Model with Autoregressive Disturbances, *Journal of Real Estate Finance and Economics*, 17:1, 99-121.
- Korbo, B.G., Rizvi, S., Ghebre, K. and Merritt, G. (2003), Location Adjustments for Agricultural Land Using the Geostatistical Capabilities of a GIS, *Assessment Journal*, Fall, 34-43.
- Lai, T-Y. and Wang, K. (1996), Comparing the Accuracy of the Minimum-Variance Grid Method to Multiple Regression in Appraised Value Estimates, *Real Estate Economics*, 24:4, 531-549.
- McCluskey, W.J. and Borst, R.A. (1998), Application of Hybrid Intelligent Appraisal Techniques Within The Field of Comparable Sales Analysis, IAAO Conference Proceedings: Today's Vision Tomorrow's Reality.
- McCluskey, W.J., Deddis, W.G., Lamont, I. and Borst, R.A. (2000), The Application of Surface Generated Interpolation Models for the Prediction of Residential Property Values, *Journal of Property Investment and Finance*, 18:2, 162-176.
- O'Connor, P.M. (1982), Making one MRA Model Behave Appropriately Across the Whole of Large County, Paper presented at the First World Congress on Computer Assisted Valuation, Cambridge, MA.
- O'Connor, P.M. and Eichenbaum, J. (1988), Location Value Response Surfaces: The Geometry of Advanced Mass Appraisal, *Property Tax Journal*, September 1988, 277-298.
- Pace, R.K. and Gilley, O.W. (1997), Using the Spatial Configuration of the Data to Improve Estimation, *Journal of Real Estate Finance and Economics*, 14:3, 333-340.
- Pace, R.K. and Gilley, O.W. (1998), Generalizing the OLS and Grid Estimators, *Real Estate Economics*, 26:2, 331-347.
- Smith, R.A. (1986), Comparing Traditional Methods for Selecting Class Intervals on Choropleth maps, *Professional Geographer*, 38:1, 62-67.
- Thibodeau, Thomas G. (2003), Marking Single-Family Property Values to Market, *Real Estate Economics*, 31:1, 1-22.
- Thompson, J.F. (2006), Comparable Sales Weighting Formulae, Cole Layer Trumble, private communication.
- Thompson, J.F. and Gordon, J.F. (1987), Constrained Regression Modeling and the Multiple Regression Analysis-Comparable Sales Approach, *Property Tax Journal*, 6:4, 35-46.

Todora, J. and Whiterell, D. (2002), Automating the Sales Comparison Approach, *Assessment Journal*, Jan/Feb, 25-33.

Underwood, W.E. and Moesch, J.R. (1982), The Second Generation of CAMA in New York State, paper presented at the First World Congress on Computer Assisted Valuation, Cambridge, MA.

Vandell, K.D. (1991), Optimal Comparables Selection and Weighting in Real Property Valuation, *Real Estate Economics*, 19:2, 213-239.

Ward, R.D., Guilford, J., Jones, B., Pratt, D. and German, J.C. (2002), Piecing Together Location: Three Studies by the Lucas County Research and Development Staff, *Assessment Journal*, Sept/Oct., 15-48.

Ward, R.D., Weaver, J.R., and German, J.C., (1999), Improving CAMA Models Using Geographic Information Systems Response Surface Analysis Location Factors, *Assessment Journal*, Jan/Feb, 30-38.

ERRATUM

In PRPRJ 12(4), the full list of authors was not shown in one paper.

Correct details are:

“Qualitative methodology and grounded theory in property research”

Deborah Levy and Christina Lee
University of Auckland